

SURWEY Study of Solriamfetol: Initiation, Titration, Safety, Efficacy, and Follow-Up Experience for Patients With OSA in Germany

Yaroslav Winter, MD¹; Geert Mayer, MD²; Sylvia Kotterba, MD³; Heike Benes, MD⁴; Lothar Burghaus, MD⁵; Iresha Abeynayake, MPH⁶; Pamela Romero-Cruz⁶; Samantha Floam, DMD⁶; Hannah Kwak⁶; Gregory Parks, PhD⁶; Ulf Kallweit, MD⁷

¹Mainz Comprehensive Epilepsy and Sleep Medicine Center, Department of Neurology, Johannes Gutenberg-University, Mainz, Germany; ²Hephata Klinik, Schwalmstadt, Germany and Philipps University Marburg, Marburg, Germany; ³Klinikum Leer gGmbH, Leer, Lower Saxony, Germany; ⁴Somni bene GmbH Institut für Medizinische Forschung und Schlafmedizin Schwerin GmbH, Schwerin, Germany; ⁵Department of Neurology, Heilig Geist-Hospital, Cologne, Germany; ⁶Axsome Therapeutics, New York, USA; ⁷Center for Biomedical Education and Research, University Witten/Herdecke, Witten, Germany

Introduction

- Excessive daytime sleepiness (EDS) is a common symptom of obstructive sleep apnea (OSA) that can persist in patients despite positive airway pressure (PAP) therapy, and may pose challenges to clinical management of symptoms¹⁻⁴
- Solriamfetol (Sunosi®) is a dopamine and norepinephrine reuptake inhibitor approved for use in adults in the European Union and United States (US) for the treatment of EDS associated with OSA (37.5–150 mg/day)^{5,6}
 - Solriamfetol has been shown to activate trace amine-associated receptor 1 (TAAR1), a potential target to improve cognitive functions^{7,8}
- There is limited real-world evidence on how physicians initiate solriamfetol and subsequent treatment outcomes in patients with OSA and associated EDS⁹; such data may help clinicians optimize patient care

Objective

- This real-world study characterizes dosing and titration strategies among European physicians initiating solriamfetol, as well as treatment outcomes following initiation in patients with EDS and OSA

Methods

- SUNOSI Real World Experience StudY (SURWEY)** was a retrospective chart review among physicians in Germany who have prescribed solriamfetol to patients with EDS associated with narcolepsy or OSA
- The present analysis focuses on data from 83 patients with OSA from Germany
 - Eligible patients were ≥18 years of age, had a diagnosis of EDS due to OSA, had reached a stable maintenance dose of solriamfetol, and completed ≥6 weeks of treatment; patients who received solriamfetol during a clinical trial or early access program were excluded
- Solriamfetol initiation strategies: **changeover** (switched/switching from existing EDS medication[s]), **add-on** (added/adding to current EDS medication[s]), and **new-to-therapy** (no current/previous EDS medication)
- Data related to solriamfetol dosing/titration, comorbidities, Epworth Sleepiness Scale (ESS) scores, patient- and physician-reported improvement in EDS, duration of solriamfetol effects, and adverse events were summarized descriptively

Results


Table 1. Baseline Demographic and Clinical Characteristics

	Changeover (n=9)	Add-on (n=12)	New-to-therapy (n=62)	Overall (N=83)
Age, mean (SD), years	44 (10)	40 (12)	51 (14)	49 (14)
Sex (female), n (%)	2 (22)	5 (42)	22 (36)	29 (35)
BMI, mean (SD), kg/m ²	33.5 (4.0)	30.7 (5.8)	32.3 (6.2)	32.2 (6.0)
ESS score, mean (SD)	16.6 (2.0)	16.3 (3.8)	15.9 (3.2)	16.0 (3.2)
Any comorbidities, n (%)	9 (100)	11 (92)	56 (90)	76 (92)
Obesity	7 (78)	6 (55)	31 (55)	44 (58)
Hypertension	4 (44)	4 (36)	29 (52)	37 (49)
Anxiety/depression	2 (22)	4 (36)	17 (30)	23 (30)
Diabetes type 2	3 (33)	2 (18)	15 (27)	20 (26)
Hyperlipidemia	5 (56)	3 (27)	10 (18)	18 (24)
Migraine headache	1 (11)	2 (18)	9 (16)	12 (16)
Congestive heart failure	0	0	8 (14)	8 (11)
Coronary artery disease	1 (11)	1 (9)	5 (9)	7 (9)
Arrhythmia	0	0	6 (11)	6 (8)
Fibromyalgia	0	1 (9)	4 (7)	5 (7)
ADHD	2 (22)	0	0	2 (3)
Other	3 (33)	5 (46)	28 (50)	36 (47)

ADHD, attention deficit hyperactivity disorder; BMI, body mass index; ESS, Epworth Sleepiness Scale; SD, standard deviation.


- New-to-therapy was the most common initiation strategy (75%), followed by add-on (14%) and changeover (11%)
- Obesity (58%) was the most common comorbidity
- Overall, 73 (88%) patients used PAP therapy; 60 (72%) patients used PAP for ≥4 hours/night and 66 (80%) patients used PAP on ≥4 nights/week
- The most commonly used concomitant medication for EDS was pitolisant

Figure 1. The Most Common Starting Dose of Solriamfetol Was 37.5 mg/day



- All 9 (100%) patients in the changeover group switched to solriamfetol due to lack of efficacy of other EDS medications
- Solriamfetol was titrated in 53 (64%) patients, most of whom (57%) completed titration within 2 weeks


Figure 2. Mean Decrease in ESS Scores^a With Solriamfetol Was ≥5 Points, Indicating Clinically Meaningful Improvement in EDS



ESS, Epworth Sleepiness Scale; SD, standard deviation. ^aScale range, 0–24; ESS scores >10 indicate EDS. ^bMean change from baseline.


- Mean improvements in ESS in changeover, add-on, and new-to-therapy groups were substantially greater than the minimum clinically important difference of 2–3 points¹¹

Figure 3. Patients and Physicians Perceived Improvements in Symptoms in Approximately 90% of Patients



EDS, excessive daytime sleepiness. ^aScale ranged from "strongly worsened" to "strongly improved"; no patients or physicians reported perceived EDS as "slightly worsened" or "strongly worsened."

Figure 4. Most Patients Reported That the Effects of Solriamfetol Lasted ≥8 Hours



- Overall, 75 (91%) patients reported no change in nighttime sleep quality with solriamfetol

Table 2. Treatment-Emergent Adverse Events^a

n (%)	Changeover (n=9)	Add-on (n=12)	New-to-therapy (n=62)	Overall (N=83)
Any TEAE	2 (22)	7 (58)	18 (29)	27 (33)
Headache	1 (11)	1 (8)	5 (8)	7 (8)
Decreased appetite	1 (11)	1 (8)	4 (6)	6 (7)
Insomnia	0	0	5 (8)	5 (6)
Feeling jittery	0	2 (17)	2 (3)	4 (5)
Irritability	0	1 (8)	3 (5)	4 (5)
Dizziness	0	2 (17)	1 (2)	3 (4)
Abdominal pain	0	0	2 (3)	2 (2)
Other	0	1 (8)	2 (3)	3 (4)

TEAE, treatment-emergent adverse event. ^aReported by ≥2 participants.

- Adverse events were consistent with those previously reported in clinical trials of solriamfetol in participants with OSA^{12,13}

Conclusions

- In the SURWEY study of solriamfetol in patients with OSA, solriamfetol was typically initiated at 37.5 mg/day and most patients were new-to-therapy; titration was common
- Overall, mean ESS scores improved by 5.4 points, and mean improvement in all groups was substantially greater than the minimum clinically important difference of 2–3 points¹¹
 - This is consistent with improvements in sleepiness reported in clinical trials for solriamfetol^{12,13}
 - Both patients and physicians perceived improvement in EDS for the vast majority of patients, regardless of initiation strategies
 - The majority of patients reported the effects of solriamfetol lasted at least 8 hours
- Common adverse events were consistent with those previously reported for solriamfetol^{12,13}

References: 1. Gasa M, et al. *J Sleep Res.* 2013;22(4):389-97. 2. Pepin JL, et al. *Eur Respir J.* 2009;33(5):1062-7. 3. Bontsignore MR, et al. *Front Neurol.* 2021;12:690008. 4. Lal C, et al. *Ann Am Thorac Soc.* 2021;18(5):757-68. 5. Sunosi® (solriamfetol) tablets Prescribing Information. New York, NY, USA: Axsome Therapeutics, Inc.; 2022. 6. Sunosi Medicine Overview. 2020. https://www.ema.europa.eu/en/documents/overview/sunosi-spar-medicine-overview_en.pdf. Accessed April 26, 2023. 7. Alnefeesi Y, et al. *Neurosci Biobehav Rev.* 2021;131:192-210. 8. Gursahani H, et al. *Sleep.* 2022;45(suppl 1):A329. 9. Winter Y, et al. *Sleep Med.* 2023;103:138-43. 10. Johns MW. *Sleep.* 1991;14(6):540-5. 11. Patel S, et al. *Am J Respir Crit Care Med.* 2018;197(7):961-3. 12. Schweitzer PK, et al. *Am J Respir Crit Care Med.* 2019;199(11):1421-31. 13. Malhotra A, et al. *Sleep.* 2020;43(2):zsz220.

Support and Acknowledgments: This study was supported by Axsome Therapeutics, Inc., Jazz Pharmaceuticals, and Pharmaviva. The development of this poster was supported by Axsome Therapeutics, Inc. On Feb. 22, 2023, Axsome Therapeutics entered into an exclusive license agreement with Pharmaviva to commercialize and further develop Sunosi® (solriamfetol) in Europe and certain countries in the Middle East and North Africa (MENA). Under the direction of the authors, Jacob Huffman, PhD, of Peloton Advantage, LLC, an OPEN Health company, provided medical writing and editorial support for this poster, which was funded by Axsome Therapeutics, Inc.

Disclosures: Y Winter has received consulting and travel support from Axsome Therapeutics and honoraria for educational presentations and consultations from Jazz Pharmaceuticals. G Mayer has no disclosures relevant to this presentation. S Kotterba, H Benes, and L Burghaus have nothing to disclose. I Abeynayake, P Romero-Cruz, S Floam, and G Parks are employees of Axsome Therapeutics. H Kwak is a former employee of Axsome Therapeutics. U Kallweit is on the advisory board at, is a consultant to, and has accepted grants/research support from Jazz Pharmaceuticals.



Scan QR code or access <https://www.axsomecongresshub.com/VASM> 2023 to view or download a PDF of this poster or access additional information and other Axsome Therapeutics presentations at VASM 2023.